

# Probability Seminar

Organizer: Christian Gromoll & Tai Melcher

Wednesday, 3:30–4:30pm, Kerchhof 317

---

Jan 17      **Adrian P. C. Lim**, Cornell

*Path integral quantization*

A typical path integral on a manifold  $M$ , is an informal expression of the form

$$\frac{1}{Z} \int_{\sigma \in H(M)} f(\sigma) e^{-E(\sigma)} \mathcal{D}\sigma, \quad (1)$$

where  $H(M)$  is a space of paths in  $M$  with energy  $E(\sigma) < \infty$ ,  $f$  is a real valued function on  $H(M)$ ,  $\mathcal{D}\sigma$  is a “Lebesgue measure” and  $Z$  is a normalization constant. The use of path integrals for “quantizing” classical mechanical systems (whose classical energy is  $E$ ) started with Feynman in [2] with very early beginnings being traced back to Dirac [1]. In this talk, I will give several rigorous definitions to Equation (1), by reviewing work done by Driver and Andersson and recently by me. The idea is to approximate  $H(M)$  by finite dimensional subspaces consisting of broken geodesics and then to pass to the limit of finer and finer approximations.

[1] P. A. M. Dirac, *Physikalische Zeitschrift der Sowjetunion* **3** (1933), 64.

[2] R. P. Feynman, *Space-time approach to non-relativistic quantum mechanics*, *Rev. Modern Physics* **20** (1948), 367–387.